

TRIANGULATED CATEGORIES OF EXTENSIONS AND THE SECOND ISOMORPHISM THEOREM FOR TRIANGULATED CATEGORIES

PETER JØRGENSEN AND KIRIKO KATO

ABSTRACT. Let \mathcal{T} be a triangulated category with triangulated subcategories \mathcal{X} and \mathcal{Y} . We show that the subcategory of extensions $\mathcal{X} * \mathcal{Y}$ is triangulated if and only if $\mathcal{Y} * \mathcal{X} \subseteq \mathcal{X} * \mathcal{Y}$.

In this situation, we show the following analogue of the Second Isomorphism Theorem: $(\mathcal{X} * \mathcal{Y})/\mathcal{X} \simeq \mathcal{Y}/(\mathcal{X} \cap \mathcal{Y})$ and $(\mathcal{X} * \mathcal{Y})/\mathcal{Y} \simeq \mathcal{X}/(\mathcal{X} \cap \mathcal{Y})$.

This follows from the existence of a stable t -structure $(\frac{\mathcal{X}}{\mathcal{X} \cap \mathcal{Y}}, \frac{\mathcal{Y}}{\mathcal{X} \cap \mathcal{Y}})$ in $(\mathcal{X} * \mathcal{Y})/(\mathcal{X} \cap \mathcal{Y})$. We use the machinery to give a recipe for constructing triangles of recollements and recover some triangles of recollements from the literature.

0. INTRODUCTION

Let \mathcal{T} be a triangulated category. If \mathcal{X} and \mathcal{Y} are full subcategories of \mathcal{T} , then the subcategory of extensions $\mathcal{X} * \mathcal{Y}$ is the full subcategory of all objects e for which there is a distinguished triangle $x \rightarrow e \rightarrow y$ with $x \in \mathcal{X}$, $y \in \mathcal{Y}$. Subcategories of extensions have been of interest to a number of authors recently, see [2], [5], [7], [13].

Our first main result is a criterion for such a subcategory to be triangulated.

Theorem A. *Let \mathcal{X} , \mathcal{Y} be triangulated subcategories of \mathcal{T} . Then $\mathcal{X} * \mathcal{Y}$ is a triangulated subcategory of $\mathcal{T} \Leftrightarrow \mathcal{Y} * \mathcal{X} \subseteq \mathcal{X} * \mathcal{Y}$.*

This is a triangulated analogue of [13, thm. 3.2]. Our second main result is the following analogue of the Second Isomorphism Theorem.

Theorem B. *Assume that \mathcal{X} , \mathcal{Y} , and $\mathcal{X} * \mathcal{Y}$ are triangulated subcategories of \mathcal{T} . Then $(\mathcal{X} * \mathcal{Y})/\mathcal{X} \simeq \mathcal{Y}/(\mathcal{X} \cap \mathcal{Y})$ and $(\mathcal{X} * \mathcal{Y})/\mathcal{Y} \simeq \mathcal{X}/(\mathcal{X} \cap \mathcal{Y})$.*

This is a consequence of the following result.

Theorem C. *Assume that \mathcal{X} , \mathcal{Y} , and $\mathcal{X} * \mathcal{Y}$ are triangulated subcategories of \mathcal{T} . Then $(\frac{\mathcal{X}}{\mathcal{X} \cap \mathcal{Y}}, \frac{\mathcal{Y}}{\mathcal{X} \cap \mathcal{Y}})$ is a stable t -structure in $(\mathcal{X} * \mathcal{Y})/(\mathcal{X} \cap \mathcal{Y})$.*

To see that this implies Theorem B, recall that a pair of triangulated subcategories $(\mathcal{U}, \mathcal{V})$ of \mathcal{T} is called a stable t -structure if $\mathcal{U} * \mathcal{V} = \mathcal{T}$ and $\text{Hom}_{\mathcal{T}}(\mathcal{U}, \mathcal{V}) = 0$, see [8, def. 9.14]. This induces equivalences of triangulated categories $\mathcal{T}/\mathcal{U} \simeq \mathcal{V}$ and $\mathcal{T}/\mathcal{V} \simeq \mathcal{U}$ by [8, prop. 9.16 and rmk. 9.17], so Theorem C induces

$$\frac{(\mathcal{X} * \mathcal{Y})/(\mathcal{X} \cap \mathcal{Y})}{\mathcal{X}/(\mathcal{X} \cap \mathcal{Y})} \simeq \frac{\mathcal{Y}}{\mathcal{X} \cap \mathcal{Y}},$$

2010 *Mathematics Subject Classification.* 18E30, 18E35, 18G35.

Key words and phrases. Homotopy category, quotient category, recollement, Second Isomorphism Theorem, stable t -structure, triangle of recollements.

that is $(X * Y) / X \simeq Y / (X \cap Y)$, see [12, prop. 2.3.1(c)]. The second equivalence of Theorem B also follows.

In Theorems B and C, we have made no comment about the important set theoretical question of whether the quotient categories have small Hom sets. However, permitting large Hom sets is not in itself a problem, cf. Neeman's treatment in [10, sec. 2.2], and if we do so, then the theorems are correct as stated. See also Remark 1.2.

Our third main result is that, under stronger assumptions, a pair of stable t-structures induces a so-called triangle of recollements in the sense of [4, def. 0.4], that is, a triple of stable t-structures (U, V) , (V, W) , (W, U) . Triangles of recollements were introduced in [4, def. 0.3] and have a very high degree of symmetry; for instance, $U \simeq V \simeq W \simeq T/U \simeq T/V \simeq T/W$. They have applications to the construction of triangle equivalences, see [4, prop. 1.16].

Theorem D. *Let (X, Y) and (Y, Z) be stable t-structures in \mathcal{T} with $Z * X = \mathcal{T}$, and let $Q : \mathcal{T} \rightarrow \mathcal{T}/(X \cap Z)$ be the quotient functor.*

Then $\mathcal{T}/(X \cap Z)$ has small Hom sets and has the stable t-structures

$$(Q(X), Q(Y)) , (Q(Y), Q(Z)) , (Q(Z), Q(X)).$$

The paper is organized as follows: Section 1 proves Theorems A, C, and D. Section 2 uses Theorem D to recover two triangles of recollements in homotopy categories known from [4, thm. 5.8] and [7, thm. 2.10].

Throughout, \mathcal{T} is a triangulated category with small Hom sets. All subcategories are full and closed under isomorphisms.

1. PROOFS OF THEOREMS A, C, AND D

Proof of Theorem A.

\Rightarrow is clear.

\Leftarrow : It is clear that $X * Y$ is closed under (de)suspension, so it is enough to show that it is closed under mapping cones.

Let $e_1 \xrightarrow{f} e_2$ be a morphism in $X * Y$. There are distinguished triangles $x_1 \xrightarrow{g} e_1 \rightarrow y_1$ and $x_2 \rightarrow e_2 \xrightarrow{h} y_2$ with $x_i \in X$ and $y_i \in Y$. The octahedral axiom gives the commutative diagram in Figure 1 with distinguished triangles as rows and columns. The mapping cone of a morphism a is denoted by $C(a)$. The distinguished triangle in the last row shows $C(hfg) \in Y * X \subseteq X * Y$, and the distinguished triangle in the last column then shows $C(fg) \in X * (X * Y) = (X * X) * Y = X * Y$. We used associativity of $*$; see [2, sec. 2.2].

Another application of the octahedral axiom gives the diagram in Figure 2. The last column shows $C(f) \in (X * Y) * Y = X * (Y * Y) = X * Y$ as desired. \square

The following consequence of Theorem A is well known, but we have been unable to locate a reference.

Corollary 1.1. *Let X, Y be triangulated subcategories of \mathcal{T} . If $\text{Hom}_{\mathcal{T}}(X, Y) = 0$ then $X * Y$ is a triangulated subcategory of \mathcal{T} .*

$$\begin{array}{ccccc}
0 & \longrightarrow & x_2 & \xlongequal{\quad} & x_2 \\
\downarrow & & \downarrow & & \downarrow \\
x_1 & \xrightarrow{fg} & e_2 & \longrightarrow & C(fg) \\
\parallel & & \downarrow h & & \downarrow \\
x_1 & \xrightarrow{hfg} & y_2 & \longrightarrow & C(hfg)
\end{array}$$

FIGURE 1. The octahedral axiom applied to fg and h

$$\begin{array}{ccccc}
x_1 & \xrightarrow{fg} & e_2 & \longrightarrow & C(fg) \\
\downarrow g & & \parallel & & \downarrow \\
e_1 & \xrightarrow{f} & e_2 & \longrightarrow & C(f) \\
\downarrow & & \downarrow & & \downarrow \\
y_1 & \longrightarrow & 0 & \longrightarrow & \Sigma y_1
\end{array}$$

FIGURE 2. The octahedral axiom applied to f and g

Proof. For $e \in Y * X$, there is a distinguished triangle $y \rightarrow e \rightarrow x \rightarrow \Sigma y$ with $x \in X$, $y \in Y$. The last morphism is 0 so $e \cong y \oplus x$ whence $e \in X * Y$ is clear. \square

Remark 1.2. If U is a triangulated subcategory of T , then we can consider the quotient category T/U . There is a quotient functor $Q : T \rightarrow T/U$. We sometimes write $Q(T)$ instead of T/U . In particular, the homomorphism functor in the quotient category is denoted $\text{Hom}_{Q(T)}(-, -)$.

Note that T/U may not have small Hom sets, but this is not in itself a problem, see [10, sec. 2.2]. In particular, we can permit large Hom sets in the situation of the following lemma and of Theorems B and C.

On the other hand, in the situation of Theorem D, we will show that the quotient $T/(X \cap Z)$ does have small Hom sets, as formulated in the theorem. This is specific to Theorem D and fails for general quotients of the form $T/(X \cap Z)$, for instance if $X = T$ and the quotient T/Z fails to have small Hom sets.

Lemma 1.3. *Assume that X , Y , and $X * Y$ are triangulated subcategories of T . Let $Q : T \rightarrow T/(X \cap Y)$ be the quotient functor.*

- (i) $\text{Hom}_{Q(T)}(Q(X), Q(Y)) = 0$.
- (ii) $Q(X) * Q(Y) = Q(X * Y)$.
- (iii) $Q(X) * Q(Y) = Q(T) \Leftrightarrow X * Y = T$.

$$\begin{array}{ccccc}
\Sigma^{-1}x' & \longrightarrow & 0 & \longrightarrow & x' \\
\downarrow & & \downarrow & & \downarrow \\
x & \xrightarrow{f} & t & \longrightarrow & C(f) \\
\downarrow & & \parallel & & \downarrow \\
w & \longrightarrow & t & \longrightarrow & y'
\end{array}$$

FIGURE 3. The octahedral axiom applied to $x \rightarrow C(f) \rightarrow v'$

Proof. (i) Given $x \in X$ and $y \in Y$, let $Qx \rightarrow Qy$ be a morphism in $Q(T)$. It is represented by a “roof”

$$\begin{array}{ccc}
& t & \\
f \nearrow & & \nwarrow g \\
x & & y
\end{array}$$

with $C(g) \in X \cap Y$. Since $y \in Y$, we have $t \in Y$.

Note that $Qx \rightarrow Qy$ is the same as $Qx \xrightarrow{Qf} Qt$ up to isomorphism, so it is enough to show $Qf = 0$. Since $X * Y$ is triangulated and contains x and t , the cone $C(f)$ is also in $X * Y$, so there is a distinguished triangle $x' \rightarrow C(f) \rightarrow y'$ in T with $x' \in X$, $y' \in Y$. The octahedral axiom gives the diagram in Figure 3. The first column shows $w \in X$ and the last row shows $w \in Y$, so we have factored f through $w \in X \cap Y$. This shows $Qf = 0$.

(ii) “ \supseteq ” is clear. “ \subseteq ”: The objects of $Q(T) = T/(X \cap Y)$ have the form Qt for $t \in T$. The condition $Qt \in Q(X) * Q(Y)$ means that there is a distinguished triangle $Qx \rightarrow Qt \rightarrow Qy$ in $Q(T)$ with $x \in X$, $y \in Y$.

Such a triangle is isomorphic to a triangle $Qa \rightarrow Qb \rightarrow Qc$ obtained by applying Q to a distinguished triangle $a \rightarrow b \rightarrow c$ from T ; see [8, def. 8.4]. There is a “roof” representing the isomorphism $Qx \cong Qa$.

$$\begin{array}{ccc}
& a' & \\
f \nearrow & & \nwarrow g \\
x & & a
\end{array}$$

Since the roof represents an isomorphism, we have $C(f), C(g) \in X \cap Y$. Since $x, C(f) \in X$, we have $a' \in X$. Since $a', C(g) \in X$, we have $a \in X$. A similar argument shows $c \in Y$. But then $b \in X * Y$ whence $Qt \cong Qb \in Q(X * Y)$.

(iii) “ \Leftarrow ” is clear. “ \Rightarrow ”: By part (ii), we need to see that $Q(X * Y) = Q(T)$ implies $X * Y = T$. But it is easy to show, using arguments similar to the ones above, that if U is any triangulated subcategory with $X \cap Y \subseteq U \subseteq T$ and $Q(U) = Q(T)$, then $U = T$. \square

Proof of Theorem C. This is immediate from Lemma 1.3(i+ii). \square

Remark 1.4. Yoshizawa gives the following example in [13, cor. 3.3]: If R is a commutative noetherian ring and S is a Serre subcategory of $\text{Mod } R$, then $(\text{mod } R) * S$ is a Serre subcategory of $\text{Mod } R$.

One might suspect a triangulated analogue to say that if \mathbb{T} is compactly generated and \mathbb{U} is a triangulated subcategory of \mathbb{T} , then so is $\mathbb{T}^c * \mathbb{U}$ where \mathbb{T}^c denotes the triangulated subcategory of compact objects. See [9, defs. 1.6 and 1.7]. However, this is false:

Set $\mathbb{T} = \mathbb{D}(\mathbb{Z})$ and $\mathbb{U} = \mathbb{D}(\mathbb{Q})$. Then \mathbb{T} is compactly generated by $\{\Sigma^i \mathbb{Z} \mid i \in \mathbb{Z}\}$. There is a homological epimorphism of rings $\mathbb{Z} \rightarrow \mathbb{Q}$ which induces an embedding of triangulated categories $\mathbb{U} \hookrightarrow \mathbb{T}$, see [3, def. 4.5] and [11, thm. 2.4]. Since \mathbb{Q} is a field, each object of \mathbb{U} has homology modules of the form $\coprod \mathbb{Q}$. This means that viewed in \mathbb{T} , the only object of \mathbb{U} which has finitely generated homology modules is 0. Hence 0 is the only object of \mathbb{U} which is compact in \mathbb{T} , see [9, cor. 2.3]. That is, $\mathbb{T}^c \cap \mathbb{U} = 0$.

If $\mathbb{T}^c * \mathbb{U}$ were a triangulated subcategory of \mathbb{T} , then Theorem C would give that $(\mathbb{T}^c, \mathbb{U})$ was a stable t-structure in $\mathbb{T}^c * \mathbb{U}$, but this is false since the canonical map $\mathbb{Z} \rightarrow \mathbb{Q}$ is a non-zero morphism from an object of \mathbb{T}^c to an object of \mathbb{U} .

Proof of Theorem D. We have that $((\mathbb{X} \cap \mathbb{Z}) \cap \mathbb{X}, (\mathbb{X} \cap \mathbb{Z}) \cap \mathbb{Y}) = (\mathbb{X} \cap \mathbb{Z}, 0)$ is (trivially) a stable t-structure in $\mathbb{X} \cap \mathbb{Z}$. Hence $(Q(\mathbb{X}), Q(\mathbb{Y}))$ is a stable t-structure in $\mathbb{T}/(\mathbb{X} \cap \mathbb{Z})$ by [4, prop. 1.5]. The second stable t-structure in the theorem is obtained analogously and the third one comes from Theorem C.

To show that $\mathbb{T}/(\mathbb{X} \cap \mathbb{Z})$ has small Hom sets, let $t, t' \in \mathbb{T}$ be given. There are distinguished triangles $y \rightarrow t \rightarrow z$ and $x' \rightarrow t' \rightarrow y'$ in \mathbb{T} with $x' \in \mathbb{X}$, $y, y' \in \mathbb{Y}$, $z \in \mathbb{Z}$. The first one gives an exact sequence

$$\mathrm{Hom}_{Q(\mathbb{T})}(Qz, Qt') \rightarrow \mathrm{Hom}_{Q(\mathbb{T})}(Qt, Qt') \rightarrow \mathrm{Hom}_{Q(\mathbb{T})}(Qy, Qt')$$

so it is enough to check that the two outer Homs are small sets. The second distinguished triangle gives similar exact sequences which show that it is enough to check that the following Homs are small sets.

$$\begin{aligned} & \mathrm{Hom}_{Q(\mathbb{T})}(Qz, Qx') \quad , \quad \mathrm{Hom}_{Q(\mathbb{T})}(Qz, Qy'), \\ & \mathrm{Hom}_{Q(\mathbb{T})}(Qy, Qx') \quad , \quad \mathrm{Hom}_{Q(\mathbb{T})}(Qy, Qy'). \end{aligned}$$

The first of these is 0 since $(Q(\mathbb{Z}), Q(\mathbb{X}))$ is a stable t-structure. Inspection of the other three shows that it is enough to check that Homs out of and into objects of $Q(\mathbb{Y})$ are small sets.

Let us check the first of these statements, as the second has a similar proof. So let $\tilde{y} \in \mathbb{Y}$ and $t'' \in \mathbb{T}$ be given. There is a distinguished triangle $y'' \rightarrow t'' \rightarrow z''$ with $y'' \in \mathbb{Y}$, $z'' \in \mathbb{Z}$. It induces a distinguished triangle $Qy'' \rightarrow Qt'' \rightarrow Qz''$ in $\mathbb{T}/(\mathbb{X} \cap \mathbb{Z})$. We have shown that $(Q(\mathbb{Y}), Q(\mathbb{Z}))$ is a stable t-structure, so $\mathrm{Hom}_{Q(\mathbb{T})}(Q\tilde{y}, \Sigma^* Qz'') = 0$. So the distinguished triangle induces a bijection

$$\mathrm{Hom}_{Q(\mathbb{T})}(Q\tilde{y}, Qy'') \rightarrow \mathrm{Hom}_{Q(\mathbb{T})}(Q\tilde{y}, Qt'').$$

It follows from [4, prop. 1.5] that $\mathrm{Hom}_{Q(\mathbb{T})}(Q\tilde{y}, Qy'') \cong \mathrm{Hom}_{\mathbb{T}}(\tilde{y}, y'')$. But $\mathrm{Hom}_{\mathbb{T}}(\tilde{y}, y'')$ is a small set by assumption, so it follows that $\mathrm{Hom}_{Q(\mathbb{T})}(Q\tilde{y}, Qt'')$ is a small set. \square

The following propositions provide pairs of stable t-structures to which Theorem D (potentially) applies.

Proposition 1.5. *Let (\mathbb{X}, \mathbb{Y}) and (\mathbb{Y}, \mathbb{Z}) be stable t-structures in \mathbb{T} with $\mathbb{Z} * \mathbb{X}$ triangulated.*

Then

$$(\mathbb{X}, \mathbb{Y} \cap (\mathbb{Z} * \mathbb{X})) \quad , \quad (\mathbb{Y} \cap (\mathbb{Z} * \mathbb{X}), \mathbb{Z})$$

*are stable t-structures in $\mathbb{Z} * \mathbb{X}$.*

$$\begin{array}{ccccc}
0 & \longrightarrow & z & \xlongequal{\quad} & z \\
\downarrow & & \downarrow & & \downarrow \\
x' & \longrightarrow & m & \longrightarrow & y' \\
\parallel & & \downarrow & & \downarrow \\
x' & \longrightarrow & x & \longrightarrow & x''
\end{array}$$

FIGURE 4. The octahedral axiom applied to $x' \rightarrow m \rightarrow x$

Proof. Let $m \in Z * X$ be given. There is a distinguished triangle $z \rightarrow m \rightarrow x$ with $z \in Z$, $x \in X$. Let $x' \rightarrow m \rightarrow y'$ be a distinguished triangle with $x' \in X$, $y' \in Y$. The octahedral axiom gives Figure 4. Since $x', x \in X$ we have $x'' \in X$ so the last column shows $y' \in Z * X$. This shows that $(X, Y \cap (Z * X))$ is a stable t-structure in $Z * X$ and a similar argument handles $(Y \cap (Z * X), Z)$. \square

If X is a subcategory of \mathbb{T} , then we write

$$X^\perp = \{t \in \mathbb{T} \mid \text{Hom}_{\mathbb{T}}(X, t) = 0\}, \quad {}^\perp X = \{t \in \mathbb{T} \mid \text{Hom}_{\mathbb{T}}(t, X) = 0\}.$$

Proposition 1.6. *Let X be a triangulated subcategory of \mathbb{T} . Set*

$$F = X * X^\perp, \quad G = {}^\perp X * X, \quad U = F \cap G.$$

Then U is a triangulated subcategory of \mathbb{T} and

$$({}^\perp X \cap F, X), \quad (X, X^\perp \cap G)$$

are stable t-structures in U .

Proof. It is immediate from Corollary 1.1 that F and G are triangulated subcategories of \mathbb{T} , so the same follows for U .

To see that $({}^\perp X \cap F, X)$ is a stable t-structure in U , observe that $\text{Hom}_U({}^\perp X \cap F, X) = 0$ is clear.

We must also show $({}^\perp X \cap F) * X = U$. The inclusion \subseteq follows because ${}^\perp X \cap F, X \subseteq U$. For the inclusion \supseteq , let $u \in U$ be given. Then $u \in G$ so there is a distinguished triangle

$$p \rightarrow u \rightarrow x \tag{1}$$

with $p \in {}^\perp X$, $x \in X$. Since $u, x \in F$ we have $p \in F$ whence $p \in {}^\perp X \cap F$, and (1) shows $u \in ({}^\perp X \cap F) * X$.

A similar argument shows that $(X, X^\perp \cap G)$ is a stable t-structure in U . \square

2. EXAMPLES

2.a. The homotopy category of projective modules. Let R be an Iwanaga-Gorenstein ring, that is, a noetherian ring which has finite injective dimension from either side as a module over itself. Let \mathbb{T} be $K_{(b)}(\text{Prj } R)$, the homotopy category of complexes of projective left- R -modules with bounded homology. Define subcategories of \mathbb{T} by

$$X = K_{(b)}^-(\text{Prj } R), \quad Y = K_{\text{ac}}(\text{Prj } R), \quad Z = K_{(b)}^+(\text{Prj } R)$$

where $\mathbf{K}_{(b)}^-(\text{Prj } R)$ is the isomorphism closure of the class of complexes P with $P^i = 0$ for $i \gg 0$ and $\mathbf{K}_{(b)}^+(\text{Prj } R)$ is defined analogously, while $\mathbf{K}_{\text{ac}}(\text{Prj } R)$ is the subcategory of acyclic (that is, exact) complexes.

Note that \mathbf{Y} is equal to $\mathbf{K}_{\text{tac}}(\text{Prj } R)$, the subcategory of totally acyclic complexes, that is, acyclic complexes which stay acyclic under the functor $\text{Hom}_R(-, Q)$ when Q is projective, see [6, cor. 5.5 and par. 5.12].

By [4, prop. 2.3(1), lem. 5.6(1), and rmk. 5.14] there are stable t-structures (\mathbf{X}, \mathbf{Y}) , (\mathbf{Y}, \mathbf{Z}) in \mathbf{T} .

If $P \in \mathbf{T}$ is given, then there is a distinguished triangle $P^{\geq 0} \rightarrow P \rightarrow P^{< 0}$ where $P^{\geq 0}$ and $P^{< 0}$ are hard truncations. Since $P^{\geq 0} \in \mathbf{Z}$ and $P^{< 0} \in \mathbf{X}$, we have $\mathbf{T} = \mathbf{Z} * \mathbf{X}$.

We can hence apply Theorem D. The intersection

$$\mathbf{X} \cap \mathbf{Z} = \mathbf{K}_{(b)}^-(\text{Prj } R) \cap \mathbf{K}_{(b)}^+(\text{Prj } R) = \mathbf{K}^b(\text{Prj } R)$$

is the isomorphism closure of the class of bounded complexes. If we use an obvious shorthand for quotient categories, Theorem D therefore provides a triangle of recollements

$$(\mathbf{K}_{(b)}^-/\mathbf{K}^b(\text{Prj } R), \mathbf{K}_{\text{ac}}(\text{Prj } R), \mathbf{K}_{(b)}^+/\mathbf{K}^b(\text{Prj } R))$$

in $\mathbf{K}_{(b)}/\mathbf{K}^b(\text{Prj } R)$. Note that $\mathbf{K}_{\text{ac}}(\text{Prj } R)$ is equivalent to its projection to $\mathbf{K}_{(b)}/\mathbf{K}^b(\text{Prj } R)$ by [4, prop. 1.5], so we can write $\mathbf{K}_{\text{ac}}(\text{Prj } R)$ instead of the projection.

This example and its finite analogue were first obtained in [4, thms. 2.8 and 5.8] and motivated the definition of triangles of recollements.

2.b. The symmetric Auslander category. Let R be a noetherian commutative ring with a dualizing complex. In [7, def. 2.1] we introduced the symmetric Auslander category $\mathbf{A}^s(R)$ as the isomorphism closure in $\mathbf{K}_{(b)}(\text{Prj } R)$ of the class of complexes P for which there exist $T, U \in \mathbf{K}_{\text{tac}}(\text{Prj } R)$ such that $P^{\leq 0} = T^{\leq 0}$ and $P^{\gg 0} = U^{\gg 0}$.

The methods of [7, sec. 2] show that there are stable t-structures

$$(\mathbf{A}(R), \mathbf{K}_{\text{tac}}(\text{Prj } R)), (\mathbf{K}_{\text{tac}}(\text{Prj } R), \mathbf{S}(\mathbf{B}(R)))$$

in $\mathbf{A}^s(R)$ with $\mathbf{S}(\mathbf{B}(R)) * \mathbf{A}(R) = \mathbf{A}^s(R)$. Here $\mathbf{A}(R)$ and $\mathbf{B}(R)$ are the Auslander and Bass categories of R , see [1, sec. 3], and \mathbf{S} is the functor introduced in [6, sec. 4.3]. It is easy to show that $\mathbf{A}(R) \cap \mathbf{S}(\mathbf{B}(R)) = \mathbf{K}^b(\text{Prj } R)$ and using Theorem D recovers the triangle of recollements

$$(\mathbf{A}(R)/\mathbf{K}^b(\text{Prj } R), \mathbf{K}_{\text{tac}}(\text{Prj } R), \mathbf{S}(\mathbf{B}(R))/\mathbf{K}^b(\text{Prj } R))$$

in $\mathbf{A}^s(R)/\mathbf{K}^b(\text{Prj } R)$ first obtained in [7, thm. 2.10]. As in the previous example, the category $\mathbf{K}_{\text{tac}}(\text{Prj } R)$ is equivalent to its projection to $\mathbf{A}^s(R)/\mathbf{K}^b(\text{Prj } R)$ so we can write $\mathbf{K}_{\text{tac}}(\text{Prj } R)$ instead of the projection.

Acknowledgement. The authors gratefully acknowledge support from JSPS Grant 22540053 and a Scheme 2 Grant from the London Mathematical Society.

REFERENCES

- [1] L. L. Avramov and H.-B. Foxby, *Ring homomorphisms and finite Gorenstein dimension*, Proc. London Math. Soc. (3) **75** (1997), 241–270.
- [2] A. Bondal and M. Van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*, Mosc. Math. J. **3** (2003), 1–36.

- [3] W. Geigle and H. Lenzing, *Perpendicular categories and applications to representations and sheaves*, J. Algebra **144** (1991), 273–343.
- [4] O. Iyama, K. Kato, and J. Miyachi, *Recollement of homotopy categories and Cohen-Macaulay modules*, J. K-Theory **8** (2011), 507–542.
- [5] O. Iyama and Y. Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, Invent. Math. **172** (2008), 117–168.
- [6] S. Iyengar and H. Krause, *Acyclicity versus total acyclicity for complexes over noetherian rings*, Doc. Math. **11** (2006), 207–240.
- [7] P. Jørgensen and K. Kato, *Symmetric Auslander and Bass categories*, Math. Proc. Cambridge Phil. Soc. **250** (2011), 227–240.
- [8] J. Miyachi, *Derived Categories with applications to representations of algebras*, Chiba University, 2000. Available at the author’s webpage: <http://www.u-gakugei.ac.jp/~miyachi/seminar.html>.
- [9] A. Neeman, *The Grothendieck duality theorem via Bousfield’s techniques and Brown Representability*, J. Amer. Math. Soc. **9** (1996), 205–236.
- [10] A. Neeman, “Triangulated categories”, Ann. of Math. Stud., Vol. 148, Princeton University Press, Princeton, 2001.
- [11] D. Pauksztello, *Homological epimorphisms of Differential Graded algebras*, Comm. Algebra **37** (2009), 2337–2350.
- [12] J.-L. Verdier, *Des catégories dérivées des catégories abéliennes*, with a preface by Luc Illusie; edited and with a note by Georges Maltsiniotis, Astérisque **239** (1996).
- [13] T. Yoshizawa, *Subcategories of extension modules by Serre subcategories*, to appear in Proc. Amer. Math. Soc. [math.AC/1011.0376v2](https://doi.org/10.1090/S0002-9947-2011-1276622-0).

SCHOOL OF MATHEMATICS AND STATISTICS, NEWCASTLE UNIVERSITY, NEWCASTLE UPON TYNE NE1 7RU, UNITED KINGDOM

E-mail address: `peter.jorgensen@ncl.ac.uk`

URL: <http://www.staff.ncl.ac.uk/peter.jorgensen>

DEPARTMENT OF MATHEMATICS AND INFORMATION SCIENCES, OSAKA PREFECTURE UNIVERSITY, OSAKA, JAPAN

E-mail address: `kiriko@mi.s.osakafu-u.ac.jp`

URL: <http://www.mi.s.osakafu-u.ac.jp/~kiriko>