

PUBLICATIONS: W. CLEGG 1987

191. Preparations, crystal structures, and molybdenum-95 nuclear magnetic resonance spectroscopic studies of triphenylphosphine-gold-tetrathiomolybdate(VI) clusters  
J. M. Charnock, S. Bristow, J. R. Nicholson, C. D. Garner and W. Clegg.  
*J. Chem. Soc. Dalton Trans.* 1987, 303–306.
192. Orthorhombic anhydrous zinc(II) propionate  
W. Clegg, I. R. Little and B. P. Straughan.  
*Acta Cryst.* 1987, **C43**, 456–457.
193. Structure and bonding of the lithium tetrahydroborate-tetramethylethylenediamine adduct (TMEDA·LiBH<sub>4</sub>)<sub>2</sub>, a centrosymmetric dimer containing doubly and triply bridging hydrogen atoms  
D. R. Armstrong, W. Clegg, H. M. Colquhoun, J. A. Daniels, R. E. Mulvey, I. R. Stephenson and K. Wade.  
*J. Chem. Soc. Chem. Commun.* 1987, 630–632.
194. Structure of tetrakis(tetraphenylphosphonium) octakis[chlorocopper(I)]bis[tetrathiotungstate(VI)]  
W. Clegg, C. D. Scattergood and C. D. Garner.  
*Acta Cryst.* 1987, **C43**, 786–787.
195. Redetermination of the structure of tetrakis(acetylacetonato)zirconium(IV)  
W. Clegg.  
*Acta Cryst.* 1987, **C43**, 789–791.
196. Structure of bis(tetramethylammonium) oxopentathiocyanatomolybdate(V)  
W. Clegg.  
*Acta Cryst.* 1987, **C43**, 791–792.
197. Structures of three glycine-bridged polymeric complexes:  
[Mn(glycine)(H<sub>2</sub>O)<sub>2</sub>Cl<sub>2</sub>], [Co(glycine)(H<sub>2</sub>O)<sub>2</sub>Cl<sub>2</sub>] and [Co(glycine)(H<sub>2</sub>O)<sub>4</sub>](NO<sub>3</sub>)<sub>2</sub>  
W. Clegg, O. M. Lacy and B. P. Straughan.  
*Acta Cryst.* 1987, **C43**, 794–797.
198. Structure of a photochromic benzoxazine derivative  
W. Clegg, N. C. Norman, J. G. Lasch and W. S. Kwak.  
*Acta Cryst.* 1987, **C43**, 804–806.
199. X-ray crystallographic and solution studies of the pentamethyldiethylenetriamine and tetramethylethylenediamine adducts of lithium diphenylphosphide  
R. E. Mulvey, K. Wade, D. R. Armstrong, G. T. Walker, R. Snaith, W. Clegg and D. Reed.  
*Polyhedron* 1987, **6**, 987–993.
200. A novel six-vertex coordination polyhedron in the crystal structure of the CuBr<sub>2</sub> complex of 1,6-bis(benzimidazol-2-yl)-2,5-dioxahexane  
W. Clegg and J. C. Lockhart.  
*Polyhedron* 1987, **6**, 1149–1151.
201. The ring-stacking principle in organolithium chemistry: its development through the isolation and crystal structures of hexameric iminolithium clusters (RR'C=NLi)<sub>6</sub> (R' = Ph, R = Bu<sup>t</sup> or Me<sub>2</sub>N; R = R' = Me<sub>2</sub>N or Bu<sup>t</sup>)  
D. R. Armstrong, D. Barr, R. Snaith, W. Clegg, R. E. Mulvey, K. Wade and D. Reed.  
*J. Chem. Soc. Dalton Trans.* 1987, 1071–1081.
202. A structural comparison of bis(quinoxaline-2,3-dithiolato)cuprate complexes  
S. Boyde, C. D. Garner and W. Clegg.  
*J. Chem. Soc. Dalton Trans.* 1987, 1083–1087.

- 203.** An unusual double C–H activation of coordinated trimethylphosphine. Synthesis, molecular structure and reactivity of  $(C_5Me_5)Ta(PMe_3)(H)_2(\eta^2-CHPMe_2)$   
T. P. Kee, V. C. Gibson and W. Clegg.  
*J. Organomet. Chem.* 1987, **325**, C14–C18.
- 204.** Preparation and crystal structure of the dicarboranylmagnesium bis(dioxane) adduct  $Mg(2-Me-1,2-C_2B_{10}H_{10})_2 \cdot 2C_4H_8O_2$   
W. Clegg, D. A. Brown, S. J. Bryan and K. Wade.  
*J. Organomet. Chem.* 1987, **325**, 39–46.
- 205.** Structure and dynamics of crowns containing the phenyldinaphthylmethane subunit (a three-bladed propeller): observations of correlated rotation of the propeller blades and certain ether segments  
J. C. Lockhart, M. B. McDonnell, W. Clegg and M. N. S. Hill.  
*J. Chem. Soc. Perkin Trans. 1* 1987, 639–649.
- 206.** Three model structural types in lithium amide chemistry: the crystal and molecular structures of dimeric  $[Ph(Me)NLi \cdot TMEDA]_2$  (TMEDA =  $Me_2NCH_2CH_2NMe_2$ ) and monomeric  $Ph(naphthyl)NLi \cdot PMDETA$  [PMDETA =  $MeN(CH_2CH_2NMe_2)_2$ ], and of  $[Ph(naphthyl)NLi \cdot TMEDA]_2$ , a ‘loose dimer’  
D. Barr, W. Clegg, R. E. Mulvey, R. Snaith and D. S. Wright.  
*J. Chem. Soc. Chem. Commun.* 1987, 716–718.
- 207.** X-ray crystal structures of some lithium aryl thiolates: monomeric  $[o-MeC_6H_4SLi \cdot (NC_5H_5)_3]$ , chain  $[PhSLi \cdot (NC_5H_5)_2]_\infty$ , and the unique folded ladder of  $[PhCH_2SLi \cdot NC_5H_5]_\infty$  with Li–S rungs  
A. J. Banister, W. Clegg and W. R. Gill.  
*J. Chem. Soc. Chem. Commun.* 1987, 850–852.
- 208.** Preparation and crystal structures of  $[PPh_4]_2[M_2(SPh)_6]$  (M = Zn or Cd)  
I. L. Abrahams, C. D. Garner and W. Clegg.  
*J. Chem. Soc. Dalton Trans.* 1987, 1577–1579.
- 209.** Structure of dichlorobis(*N*-isopropylideneaniline)palladium(II)  
W. Clegg, J. Barker and M. Kilner.  
*Acta Cryst.* 1987, **C43**, 1211–1212.
- 210.** A pentuply-bridging carbonyl group: crystal and molecular structure of a salt of the 1-oxo-2-phenyl-1,2-dicarbododecaborate(12) anion,  $[LH]^+[O(Ph)C_2B_{10}H_{10}]^-$  (L = 1,8-*N,N,N',N'*-tetramethylnaphthalenediamine)  
D. A. Brown, W. Clegg, H. M. Colquhoun, J. A. Daniels, I. R. Stephenson and K. Wade.  
*J. Chem. Soc. Chem. Commun.* 1987, 889–891.
- 211.** A new route to aryltin compounds: orthometallation of Schiff’s bases by tin(IV) halides: crystal and molecular structure of *N*-methyl-*ortho*-trichlorostannyldiphenylmethylethylamine,  $2-Cl_3SnC_6H_4C(Ph)=NMe$   
W. Clegg, C. M. J. Grievson and K. Wade.  
*J. Chem. Soc. Chem. Commun.* 1987, 969–970.
- 212.** Regioselective ring-opening and dimerisation of methyl 3,3-dimethylcyclopropenecarboxylate  
M. S. Baird, H. H. Hussain and W. Clegg.  
*J. Chem. Soc. Perkin Trans. 1* 1987, 1609–1612.
- 213.** Structure and bonding of the tris(diphenylmethylethylamino)aluminium dimer  $[Al(NCPh_2)_3]_2$ : relative strengths of bridging and terminal Al–X bonds in systems  $X_2Al(\mu_2-X)_2AlX_2$   
S. J. Bryan, W. Clegg, R. Snaith, K. Wade and E. H. Wong.  
*J. Chem. Soc. Chem. Commun.* 1987, 1223–1224.

- 214.** Structure of tetrakis( $\mu$ -1,3-benzothiazole-2-thiolato-*N,S*)-diplatinum(II) dimethylformamide solvate  
E. S. Raper, A. M. Britton, J. R. Creighton, W. Clegg, M. Hooper and M. Kubiak.  
*Acta Cryst.* 1987, **C43**, 1538–1541.
- 215.** The syntheses and crystal structures of two iminolithium adducts:  $(\text{Ph}_2\text{C}=\text{NLi}\cdot\text{NC}_5\text{H}_5)_4$ , a tetrameric pseudo-cubane formed by stacking two  $(\text{LiN})_2$  four-membered rings, and  $[\text{Bu}^t_2\text{C}=\text{NLi}\cdot\text{OP}(\text{NMe}_2)_3]_2$ , a dimer prevented sterically from stacking  
D. Barr, R. Snaith, W. Clegg, R. E. Mulvey and K. Wade.  
*J. Chem. Soc. Dalton Trans.* 1987, 2141–2147.
- 216.** Isolation of the single cubane-like Re–S–Cu–Cl cluster  $[\text{ReS}_4(\text{CuCl})_3\text{Cl}]^{2-}$   
C. D. Scattergood, C. D. Garner and W. Clegg.  
*Inorg. Chim. Acta* 1987, **132**, 161–162.
- 217.** Synthesis and X-ray crystal structure of  $[\{(\text{C}_5\text{H}_4\text{Me})\text{Fe}(\text{CO})_2\}_2\text{BiCl}]_3$ : a compound containing a planar six-membered  $\text{Bi}_3\text{Cl}_3$  ring  
W. Clegg, N. A. Compton, R. J. Errington and N. C. Norman.  
*Polyhedron* 1987, **6**, 2031–2033.
- 218.** Reactions of  $\text{S}_3\text{N}_2\text{Cl}_2$  and  $(\text{NSCl})_3$  with a new silylcyanamide reagent  $(\text{Me}_3\text{Si})_2\text{N}\cdot\text{CN}$ : synthesis and X-ray crystal structure of  $\text{SNSNS:N}\cdot\text{CN}$  and bicyclic  $\text{CClS}_3\text{N}_5$   
A. J. Banister, W. Clegg, I. B. Gorrell, Z. V. Hauptman and R. W. H. Small.  
*J. Chem. Soc. Chem. Commun.* 1987, 1611–1613.
- 219.** On the possible isomerisation pathway for triarylmethanes  $\text{Ar}_3\text{CX}$ , including propeller crowns  
W. Clegg and J. C. Lockhart.  
*J. Chem. Soc. Perkin Trans. 1* 1987, 1621–1627.
- 220.** The synthesis of some organo-molybdenum and -tungsten complexes containing bismuth  
W. Clegg, N. A. Compton, R. J. Errington and N. C. Norman.  
*J. Organomet. Chem.* 1987, **335**, C1–C5.
- 221.** Structure of di- $\mu$ -chloro-bis[trichloro(*p*-tolylimino)tungsten(VI)],  $[\{\text{WCl}_3(\text{p-CH}_3\text{C}_6\text{H}_4\text{N})\}_2(\mu\text{-Cl})_2]$   
W. Clegg and R. J. Errington.  
*Acta Cryst.* 1987, **C43**, 2223–2224.
- 222.** Synthesis and characterisation of  $[\text{VS}_4(\text{CuPPh}_3)_5(\text{CuCl})\text{Cl}_2]\cdot\text{CH}_2\text{Cl}_2$ ; a compound containing a tetrahedral  $\text{V}^{\text{V}}\text{S}_4$  core encapsulated by an octahedral array of copper(I) atoms  
C. D. Scattergood, P. G. Bonney, J. M. Slater, C. D. Garner and W. Clegg.  
*J. Chem. Soc. Chem. Commun.* 1987, 1749–1750.