

Use and husbandry of captive European starlings (*Sturnus vulgaris*) in scientific research: a review of current practice

Lucy Asher* and Melissa Bateson†

*Epidemiology Division, Royal Veterinary College, Hawkshead Campus, Hatfield, Hertfordshire AL9 7TA, UK; †School of Biology and Psychology, Newcastle University, Newcastle upon Tyne, UK

Summary

We reviewed the use of captive European starlings (*Sturnus vulgaris*) in scientific research published between 2000 and 2004. We estimated the numbers of birds used and documented their origin and the range of husbandry regimes employed with the aim of comparing current practice with the new European guidelines for husbandry of laboratory animals. Over the five-year period, 106 primary articles report the use of an estimated total of 2490 captive starlings. The majority of birds were caught from the wild either as adults or fledglings, and only 3% were hand-reared from chicks. There was considerable variation in husbandry. In the majority of cases, standards fell below those currently recommended as best practice in the UK and cited in new European guidelines. The median volume of home cages employed was 0.42 m³ (0.13–5.1 m³, interquartile range), whereas current recommendations suggest a minimum of 1.0 m³ for a singly-housed bird. The median volume of space allowed per bird was 0.13 m³/bird (0.08–1.05 m³/bird, Q1–Q3), whereas current recommendations suggest a minimum of 0.33 m³/bird. Only 27% of the articles mentioned providing any form of environmental enrichment for birds in their home cages. We recommend that more research be conducted into the welfare of starlings to inform legislation and guidelines, and thus maximize the welfare of captive animals.

Keywords Husbandry; laboratory animal welfare; legislation; starling; *Sturnus vulgaris*

The European Union (EU) is currently drafting revisions of legislation relating to the housing of laboratory animals. Major changes under consideration include mandatory minimum cage sizes, environmental enrichment and group housing, and limitations on the use of wild-caught animals in research (European Commission 2006). For commonly used laboratory species, these specifications are often evidence-based and are claimed to reflect current practice in many laboratories. However, this is not the case for many less common but nevertheless important laboratory species, in which there has been little welfare research, and for which husbandry is not standardized

between laboratories. We review the use and husbandry of European starlings (*Sturnus vulgaris*) to quantify current practice, and thus assess the impact of adherence to new animal welfare guidelines in laboratories using this species.

The European starling is arguably the most widely used passerine species in fundamental biological research. A search on ISI Web of Knowledge reveals 392 articles listing 'Sturnus vulgaris' in the topic field in the five-year period from 2000–2004 (compared with only 276 listing 'Taeniopygia guttata', the zebra finch). For many years now starlings have been an important model species in studies of behavioural ecology, ecophysiology and the neuroecology of song production and perception (e.g. Powell 1974, Tinbergen 1981, Lima 1983, Kalcelnik 1984,

Correspondence: L. Asher. Email: lasher@rvc.ac.uk

Accepted 18 May 2007

Cuthill & Hindmarsh 1985, West & King 1990, Eens *et al.* 1991, Mountjoy & Lemon 1991). The popularity of starlings stems from a number of features that make them ideal experimental animals. Starlings acquire new tasks rapidly and can readily be trained on operant schedules making them ideal subject for studies of foraging behaviour (e.g. Bateson & Kacelnik 1995). Their complex songs and propensity for vocal mimicry make them interesting subjects for studies of bird song. Although not easy to breed in captivity, wild starlings are common in both North America and Europe populating nearly 30% of the world's land surface excluding Antarctica (Feare 1984), and are easy to catch in large numbers. They adapt well to a captive environment, and due to their social nature can be housed at relatively high densities in the laboratory.

The welfare of starlings housed in the laboratory is of particular concern for three reasons. First, as non-domesticated species starlings have not been selectively bred to cope in captive environments. Secondly, starlings used in laboratory research are often caught from the wild, and have therefore experienced the loss of their natural environment. Thirdly, wild starlings travel more than 20 km daily to and from their roosting sites (Feare 1984), and in carnivores a large natural range is a risk factor for poor welfare in captivity (Clubb & Mason 2003). There is currently little data available on the welfare of captive laboratory birds (Poole & Dawkins 1999). In starlings specifically, a handful of studies have addressed the effects of lighting quality (Greenwood *et al.* 2002, 2004) and environmental enrichment (Gill 1994, Gill *et al.* 1995) on welfare, but to date there are no studies on the effects of developmental history (wild-caught versus hand-raised or captive-bred), cage size or stocking density.

Laboratory animal research in many countries falls under the Council of Europe Convention, ETS 123 and the European Union Directive 86/609/EEC for the protection of vertebrate animals used for experimental and other scientific purposes. Appendix A of ETS 123 has recently been revised with the aim of increasing standards

of welfare for laboratory animals and harmonizing practice across laboratories. Although the situation remains unresolved, it is possible that the revised Appendix A of ETS 123 may be transcribed into Annex II of European Union Directive 86/609/EEC. Also under consideration is the possibility of making elements of Annex II minimum standards that would be legally binding for all EU member states (European Commission 2006). In the revised version of ETS 123, general provisions for birds include group housing, environmental enrichment and use of captive-bred birds wherever possible (Hawkins *et al.* 2003). Although there are no specific provisions for undomesticated passerines such as the starling, readers are referred to the *Laboratory Animals'* supplement *Laboratory birds: refinements in husbandry and procedures* (Hawkins *et al.* 2001) that does provide guidelines specific to starlings. A minimum cage size of 1 m³ is recommended for a singly-housed starling, although group housing is preferable with a minimum cage size of 2 m³ for a group of two to six birds (i.e. a minimum space allowance of 0.33 m³/bird). Enrichment with natural branches, water baths, provision of cover, live invertebrate prey and a substrate of bark chippings are also recommended. Although adherence to these guidelines is not compulsory in the revised Council of Europe Convention (ETS 123), husbandry that strongly diverges from those recommended in this document is unlikely to be viewed as acceptable by the responsible authorities, at least until further research is produced. Furthermore, should the changes to Directive 86/609/EEC outlined above be agreed, general provisions could become mandatory for starlings housed in laboratories in EU countries. Therefore, all researchers in Europe will need to attend to both the Hawkins *et al.* (2001) and Hawkins (2003) documents (henceforth referred to as 'new European guidelines').

In this article, we review the use of captive European starlings in scientific research. A specific aim is to quantify current practice in the husbandry of starlings in relation to new European guidelines.

Materials and methods

Studies and extraction of information

Studies using captive European starlings (*S. vulgaris*) published between 2000 and 2004 were identified from searches on databases Agricola, Biosis, CAB, Cambridge Scientific Abstracts (BioSciences), PubMed (Medline), Web of Science and Zoological record. Searches were made between January and April 2005 using only the term 'Sturnus vulgaris' in the title/subject/keywords, and then relevant articles were selected by scanning the abstract or entire article if necessary. Articles were included in this review if they reported holding European starlings in captivity for one day or longer. A starling was deemed as captive if it was confined by a cage, aviary, room, box or any other form of restraint.

Basic information on the country of the research institution and any legislation or ethical guidelines reported, were extracted from each relevant article. We also extracted the number of male, female and total starlings used in the study, the origin of the birds, the length of time they had spent in captivity, any procedures carried out on the birds, the type of science being investigated and the fate of birds used. Details on husbandry and care included: the housing type and size, stocking density, type and availability of food and water, temperature, details of lighting and if any environmental enrichment was provided. Whether lighting was artificial or natural was noted. If lighting was artificial the hours of light and dark (lighting schedule), whether there was a gradual transition between these, the brightness, spectrum and rate of flicker, were also recorded.

Classifications, definitions and analysis

We used the database Biosis' Major Concept Terms to classify the type of science. The specific types of research conducted on captive starlings were identified by determining institutions with four or more publications in the review. Numbers of birds were calculated by summing the total numbers of birds in relevant studies. Medians (M) and interquartile ranges

(Q1–Q3) were calculated for many aspects of husbandry using the number of articles.

Dimensions were calculated for the home cage, defined as the enclosure the birds spent most time in during the experiment. The length was defined as the longer of the two horizontal dimensions and the width the shorter (the height is assumed to be reported as the vertical dimension). The space available to captive starlings was expressed in four ways: total cage volume and cage volume per bird, floor area and cage aspect ratio.

Volumes were calculated by multiplying the three cage dimensions and are always expressed in cubic metres. Home-cage volumes were divided by the maximum number of birds held in that home cage, to provide the volume per bird, which was expressed in m³/bird. Floor areas were calculated by multiplying the length and width (m²). Finally, aspect ratios were calculated to describe the shape of the profile of the cage, by dividing the length by the height. A cage with an aspect ratio <1 is taller than it is long; an aspect ratio of exactly one has a square profile; and a cage with an aspect ratio of >1 is longer than it is tall. Environmental enrichment was operationally defined as any item provided with the goal of improving the animals' welfare.

Scientific procedures were described as either invasive or non-invasive according to the UK Animals (Scientific Procedures) Act (1986). Non-invasive procedures fall into two categories: handle and restraint only, and attaching a foreign object. Non-invasive procedures were not included if they were deemed to be part of usual husbandry routines, such as trimming claws or bills.

Results

Studies: number, subject and geographical location

Information concerning the husbandry and care of captive starlings was extracted from a total of 106 research articles (see Table 1 for summary). The articles were published in 32 different journals, with the most articles sourced from *Animal Behaviour* (8 articles), *General and Comparative Endocrinology*

Table 1 Summary of the articles reviewed

	Science	Origin	Home-cage dimensions (cm)			Birds per cage	Country	Number of birds
			Length	Width	Height			
Absil <i>et al.</i> (2003a)	Neural coordination	Wild				Belgium	15	
Absil <i>et al.</i> (2003b)	Neural coordination	Wild			1	Belgium	34	
Atasver and Gumussoy (2004)	Animal care	Wild				Turkey	24	
Auger <i>et al.</i> (2002)	Sensory reception	Wild	45	75	1+	USA	16	
Bateson (2002)	Behaviour	Wild	510	200	1	UK	8	
Bautista <i>et al.</i> (2001)	Behaviour	Wild			1	UK	4	
Bee and Klump (2004)	Neural coordination	Wild				Germany	4	
Bentley (2003)	Chemical coordination	Wild	49	95	6	USA	20	
Bentley <i>et al.</i> (2000)	Chemical coordination	Wild	60	50	4-6	UK	12	
Blackwell <i>et al.</i> (2002)	Pest control	Wild	240	240	180	USA	12	
Braaten (2000)	Sensory reception	Wild			1	USA	4	
Buchanan <i>et al.</i> (2003)	Sensory reception	Wild	180	180	90	UK	48	
Butler <i>et al.</i> (2001)	Virology	Wild	61	36	41	USA	8	
Clark <i>et al.</i> (2000)	Pest control	Wild			1	USA	40	
Congiu <i>et al.</i> (2000)	Pest control	Wild			1	USA	33	
Cousillas <i>et al.</i> (2004)	Neural coordination	Wild			1	France	8	
Cuthill <i>et al.</i> (2000)	Behaviour	Wild	100	30	40	UK	6	
Dawson (2001)	Chemical coordination	Wild			14	UK	95	
Dawson (2003a)	Reproduction	Wild	200	300	400	UK	10	
Dawson (2003b)	Physiology	Wild			10	UK	16	
Dawson (2004a)	Sensory reception	Wild			4/5	UK	69	
Dawson (2004b)	Physiology	Wild	310	310	260	UK	14	
Dawson <i>et al.</i> (2000)	Reproduction	Wild			8	UK	16	
Dawson <i>et al.</i> (2002)	Chemical coordination	Wild	60	40	40	UK	126	
Dawson and Van't Hof (2002)	Chemical coordination	Wild/hand-raised			4	UK	34	
De Cock and Matthysen (2001)	Sensory reception	Wild	135	175	200	Belgium	11	
De Ridder <i>et al.</i> (2002)	Chemical coordination	Wild	700	400	250	Belgium	15	
Devereux <i>et al.</i> (2004)	Behaviour	Wild	90	70	60	UK	20	
Duffy and Ball (2002)	Evolution and adaptation	Wild	250	250	150	USA	16	
Duffy <i>et al.</i> (2000)	Chemical coordination	Wild			1-4	USA	40	
Fawcett <i>et al.</i> (2002)	Behaviour	Wild	60	66	37	UK	48	
Fernández-Juricic and Kacelnik (2004)	Behaviour	Wild	90	70	60	UK	24	
Fernández-Juricic <i>et al.</i> (2004)	Behaviour	Wild	90	70	60	UK	24	
Gentner and Hulse (2000a)	Sensory reception	Wild	150	500	200	USA	22	
Gentner and Hulse (2000b)	Sensory reception	Wild			1-5	USA	14	
Gentner and Margoliash (2003)	Neural coordination	Wild			1-6	USA	11	
Gentner <i>et al.</i> (2000)	Neural coordination	Wild			1-6	USA	20	
Gentner <i>et al.</i> (2001)	Sensory reception	Wild	200	200	150	USA	27	

Gentner <i>et al.</i> (2004)	Sensory reception	Wild						USA	17
George <i>et al.</i> (2004)	Sensory reception	Wild						France	6
Goslow <i>et al.</i> (2000)	Physiology	Wild						USA	10
Greenwood <i>et al.</i> (2002)	Animal care	Wild	130	65	120	8		UK	132
Greenwood <i>et al.</i> (2003)	Sensory reception	Wild/hand-raised	140	65	120	12		UK	8
Greenwood <i>et al.</i> (2004)	Animal care	Wild	130	65	120	12		UK	40
Gwinner <i>et al.</i> (2002)	Chemical coordination	Wild/hand-raised	400	200	300			Germany	52
Hambly <i>et al.</i> (2004)	Physiology	Wild	540	70	80	1		UK	9
Hausberger <i>et al.</i> (2000)	Neural coordination	Wild						France	4
Henry and Hausberger (2001)	Sensory reception	Wild/captive born/ hand-raised	180	1000	400			USA	17
Hile (2004)	Pest control	Wild	33	33	61	1		USA	24
Hile <i>et al.</i> (2004)	Pest control	Wild						USA	24
Hodgeson <i>et al.</i> (2001)	Virology	Wild	30	30	30	1		USA	5
Hofer and Klump (2003)	Neural coordination	Wild	35	27	52	8		Germany	8
Kacelnik and Marsh (2002)	Behaviour	Wild	120	50	53	1		UK	12
Kumar <i>et al.</i> (2000)	Chemical coordination	Wild	81	50	31	1		Germany	8
Langerman and Klump (2001)	Neural coordination	Wild	80	40	40	1		Germany	5
Lucas <i>et al.</i> (2002)	Sensory reception	Hand-raised	100	100	100	1		USA	8
Maddocks <i>et al.</i> (2001)	Sensory reception	Wild	40	30	30	8		UK	32
Maddocks <i>et al.</i> (2002a)	Animal care	Wild	70	120	50	4		UK	8
Maddocks <i>et al.</i> (2002b)	Animal care	Wild	67	120	50	8		UK	8
Maddocks <i>et al.</i> (2002c)	Sensory reception	Wild	70	90	180	8		UK	48
Marsh and Kacelnik (2002)	Behaviour	Wild	158	56	53	1		UK	16
Marsh <i>et al.</i> (2004)	Behaviour	Wild	120	60	50	1-6		UK	12
Nephew and Romero (2003)	Chemical coordination	Wild	34	38	45			USA	24
Nephew <i>et al.</i> (2003)	Reproduction	Wild	25	53	35	4-12		USA	24
Nieder and Klump (2001)	Neural coordination	Wild	32	24	25	1		Germany	8
Oberweger and Goller (2001)	Physiology	Hand-raised				1		USA	3
Olsson <i>et al.</i> (2001)	Behaviour	Wild	100	100	50	3		Sweden	12
Olsson <i>et al.</i> (2002)	Behaviour	Wild				5		Sweden	10
Pilz <i>et al.</i> (2004)	Development	Hand-raised						USA	60
Pinxten <i>et al.</i> (2000)	Chemical coordination	Wild	175	135	200	1		Belgium	18
Pinxten <i>et al.</i> (2002)	Chemical coordination	Wild	500	200	250	2-12		Belgium	35
Pinxten <i>et al.</i> (2003a)	Neural coordination	Wild	175	135	200	1-2		Belgium	24
Pinxten <i>et al.</i> (2003b)	Reproduction	Wild	175	135	200	1-4		Belgium	23
Poirier <i>et al.</i> (2004)	Sensory reception	Wild/hand-raised	40	60	52	1-5		France	23
Rafacz and Templeton (2003)	Behaviour	Wild	60	45	45	1		USA	22
Rayner <i>et al.</i> (2001)	Physiology	Wild				1		UK	1
Remage-Healey and Romero (2000)	Chemical coordination	Wild				1		USA	11
Remage-Healey and Romero (2001)	Chemical coordination	Wild				1		USA	10
Remage-Healey and Romero (2002)	Chemical coordination	Wild				2		USA	10
Riters and Alger (2004)	Sensory reception	Wild						USA	20
Riters and Teague (2003)	Chemical coordination	Wild	51	97	51			USA	24

(Continued)

Table 1 Continued

	Home-cage dimensions (cm)				Birds per cage	Country	Number of birds
	Length	Width	Height				
Riters <i>et al.</i> (2000)	600	400	250		12	USA	44
Riters <i>et al.</i> (2001)					12	USA	74
Riters <i>et al.</i> (2002)						USA	58
Romero and Remage-Healey (2000)					1	USA	25
Rothery <i>et al.</i> (2001)						UK	10
Sayre and Clark (2001)	36	23	28		1	USA	40
Schuck-Paim and Kacelnik (2002)	60	50	120		1	UK	18
Schuck-Paim <i>et al.</i> (2004)	120	60	50		1	UK	28
Seibels <i>et al.</i> (2003)	180	90	120		15	USA	61
Sockman <i>et al.</i> (2002)					4	USA	20
Sockman <i>et al.</i> (2004)					2	USA	32
Staples <i>et al.</i> (2002)				Wild/hand-raised	1	USA	10
Swaddle and Lockwood (2003)	350	120	200			USA	43
Swaddle and Pruett-Jones (2001)					1	USA	12
Swaddle and Ruff (2004)	100	60	70		1-4	USA	8
Swaddle <i>et al.</i> (2004)	90	40	40		1-3	USA	34
Talling <i>et al.</i> (2002)	710	300	250		1-6	UK	6
Templeton and Gonzalez (2004)	46	60	40		1	USA	12
Van Meir <i>et al.</i> (2004)	140	220	210		5	Belgium	10
Vasquez and Kacelnik (2000)	150	50	50		1	UK	8
Výboh <i>et al.</i> (2001)				Wild/hand-raised		Slovakia	6
Ward <i>et al.</i> (2001)	200	200	200		3-4	UK	15
Ward <i>et al.</i> (2004)	250	250	300		1	UK	4
Williams and Swaddle (2004)	49	95	51		1-7	UK	7
Young <i>et al.</i> (2001)					1-20	USA	71

(9 articles) and *Hormones and Behaviour* (7 articles). Studies spanned 11 categories of scientific research (Figure 1). Specific areas of research include: behavioural and cognitive ecology of foraging behaviour and mate choice (e.g. Maddocks *et al.* 2002a–c, Fernández-Juricic *et al.* 2004), song perception (e.g. Langemann & Klump 2001, Gentner *et al.* 2004), behavioural neuro-endocrinology of singing and the stress response (e.g. Auger *et al.* 2002, Nephew & Romero 2003, Pinxten *et al.* 2003a,b) and photoperiodic control of reproduction (e.g. Dawson 2001).

The research was produced by 32 institutions, with the vast majority being conducted in either the UK (37 articles) or the USA (45 articles), the remaining research being conducted in other European countries, Belgium (9 articles), Germany (6 articles), France (4 articles), Sweden (2 articles), Italy (1 article), Slovakia (1 article) and Turkey (1 article).

Although no articles specifically cited European legislation, 44 articles did reference some type of guidelines the study had adhered to, most commonly national legislation; seven articles mentioned UK Animals

(Scientific Procedures) Act (1986) (covering an estimated 375 birds) and 10 articles mentioned other national legislation (175 birds). Seventeen articles reported coverage by university guidelines or ethics committees (364 birds). The National Research Councils' *Guide for the Care and Use of Laboratory Animals* (1996) was cited by 16 articles (341 birds). Nine articles (322 birds) also cited the body that granted permission for capture of wild birds, and included English Nature, Scottish Natural Heritage and the US Department of Agriculture.

Bird numbers and origins

A total of 2490 starlings were used in the studies reviewed with 18, 8–29 birds (M, Q1–Q3) used per article. Number of birds used per article varied from a single bird in an experiment concerning the physiology and mechanics of flight (Rayner *et al.* 2001) to 132 birds used in Greenwood *et al.*'s (2002) study concerning preferred lighting environments. Birds were predominantly male, 742 males when compared with 346 females, although only half the articles reported information on sex.

Ninety-eight out of the total 106 articles provided some information concerning the origins of the birds used. Most studies used wild-caught birds (87 articles, 2153 birds), three articles used hand-raised birds (71 birds) and eight articles used a combination of hand-raised and wild-caught birds (165 birds). Only one article (2 birds) used captive-bred birds (Henry & Hausberger 2001).

Home cage

Of the 106 articles reviewed, 100 provided some information concerning the type of housing, but only 65 of these gave the exact cage dimensions. The enclosures varied in size and volume, ranging from 0.02 (Sayre & Clark 2001) to 720 m³ (Henry & Hausberger 2001; Figure 2). Many of the values fall within the lower range of volumes (M, Q1–Q3 = 0.42, 0.13–5.1 m³) with 65% (42 articles) falling below the 1 m³ threshold. The distribution of floor areas (Figure 3) was also positively skewed (M, Q1–Q3 = 0.75, 0.32–3.78 m²). Most enclosures were longer




Figure 1 Histogram of the types of science conducted with captive European starlings 2000–2004 for the number of articles and the number of individual birds involved. Categories are the Major Concept Terms used in the Biosis' database




Figure 2 Frequency distributions of home-cage volumes for (a) the number of articles, and (b) the number of individual birds involved. The dashed line indicates a cage volume of 1 m^3 (minimum cage size recommended in new European guidelines)

than they were tall (Figure 4), with a median aspect ratio of 0.88, 0.32–0.63 (M, Q1–Q3).

In the majority of articles birds were either housed singly (49 articles, 862 birds) or in small groups of 3–10 birds (38 articles, 1086 birds), although during the course of many studies birds experienced several different groupings (Table 2). For the 54 articles that included information on both home-cage dimensions and number of birds per cage, the volume of space per bird was 0.13, 0.08–1.05 m^3/bird (M, Q1–Q3, Figure 5).

Food

In 54 out of 71 articles that provided the information, birds received food *ad libitum* (1455 birds). The remaining 17 articles (352 birds) used some degree of food deprivation. The basic diet most fed to captive starlings was dry food (63 articles, 1589 birds). Dry




Figure 3 Frequency distributions of home-cage floor areas for (a) the number of articles, and (b) the number of individual birds involved

food included, turkey starter crumbs, Orlux pellets (Orlux, Wielsbeke-Ooigem, Belgium), Purina chick food (Purina, St Louis, USA), poultry pellets, dog food and wheat or barley. The diet of many birds was supplemented with fruit (7 articles, 193 birds), processed meat (6 articles, 179 birds), vitamin supplements (5 articles, 105 birds). Twenty-three articles reported feeding live invertebrate prey to captive starlings. Mealworms (*Tenebrio molitor*) were fed to birds in 22 articles (339 birds), six of these also provided the birds with natural forage (101 birds), and one article provided the birds with just natural forage (6 birds).

Environmental enrichment

The most commonly used forms of enrichment were nest boxes (16 articles, 468 birds) and water baths (15 articles, 306 birds). Fewer articles reported presenting birds with toys (3 articles, 88 birds), natural branches




Figure 4 Frequency distributions of home-cage aspect ratios for (a) the number of articles, and (b) the number of individual birds involved

(3 articles, 45 birds) or foraging substrate (5 articles, 58 birds). A total of 29 articles reported the use of some form of environmental enrichment.

Environmental variables

All 106 articles provided information about lighting regimes used, but 41 gave information about temperature. Temperatures

Table 2 Number of captive starlings per enclosure

	Articles	Birds
Single	27	386
Pair	2	80
Small group*	24	766
Colony†	8	349
Single + small group	13	312
Single + colony	9	164
Pair + colony	2	67
Group + colony	1	8

*A small group consists of 3–10 birds per enclosure

†A colony is more than 10 birds per enclosure




Figure 5 Frequency distributions of the amount of space per bird in the home cages for (a) the number of articles, and (b) the number of individual birds involved. The smaller dashed line indicates the new European guidelines space allowance per bird for a group-housed bird (0.33 m³/bird) and the larger dashed line the space allowance per bird for a singly-housed bird (1 m³/bird)

ranged from 12–28°C, with a median of 18 (15.75–21.50°C, Q1–Q3). A total of 95 articles used artificial lighting and although most reported the lighting schedule (83 articles; Tables 3 and 4) only 10 articles mentioned details of the lighting quality (e.g. light to dark transition, flicker rate, UV, spectrum, fluorescence, luminance).

Table 3 Number of different lighting schedules used in the articles reviewed and the number of birds this covers

Number of lighting schedules	Articles	Birds
1	60	1274
2	14	364
3	7	222
4	1	69

Table 4 Lighting schedules used and the number of birds

Lighting schedule	Articles	Birds
8L:16D	18	88
9L:15D	2	259
10L:14D	3	384
11L:13D	10	24
12L:12D	17	229
13L:11D	2	180
14L:10D	5	56
16L:8D	5	655
18L:6D	9	480
19L:5D	4	77
20L:4D	1	56
Continuous	1	69
Ambient	18	6

L=light; D=dark

Scientific procedures

Non-invasive procedures that involved attaching a foreign object included, coloured leg rings, eye rings and in one particular study a gas mask fixed to a bird's head. Handle and restraint procedures involved measuring feathers with callipers, weighing, restraint in a cloth bag for up to 30 min and restraint in a wing cuff and head holder. Little consistency over reporting these non-invasive procedures was found and frequency of handling was never comprehensively reported, e.g. some articles referred to a regular weighing scheme but few specified whether weighing was remote or involved handling.

Invasive procedures were more reliably reported. More than half (56) of the articles reviewed involved some type of procedure that would in the UK need to be covered by a Home Office licence (Figure 6).

Only 33 articles reported information about the fate of birds after involvement in research. In 26 articles (754 birds), birds were euthanized at the end of experiments either by an overdose of anaesthetic or decapitation. Birds were released upon completion of six studies (84 birds) and two articles reported birds being retained for further study (42 birds).

Discussion

Out of the 106 articles and 2490 birds in this review, 61 articles (1383 birds) were conducted in EU countries that will be subject to new European guidelines for starling




Figure 6 Histogram of the types of invasive experimental procedures conducted on European starlings for the number of articles and the number of individual birds involved. *Dosing includes the injection of any type of substance into the bird

husbandry. A comparison of these new guidelines with current practice reveals that birds were only group-housed for the duration of experiments in 43% of articles, only 27% of articles referred to the use of environmental enrichment and only two birds in the entire review were captive-bred. Furthermore, the species-specific recommendations represent a departure from common practice, with the 1 m³ minimum cage volume not observed in 42 articles (1099 birds). The median volume of home cages was 0.42 m³, and the median space allowance per bird was 0.13 m³/bird, compared with the recommended 0.33 m³/bird. The advised enrichment with live prey was used in 23 articles (345 birds), water baths in 15 articles (306 birds), foraging substrate in five articles (58 birds), natural branches in three articles (45 birds) and the provision of cover was not mentioned in any article reviewed.

It is difficult to interpret how this divergence from recommended guidelines may impact the welfare of the birds involved, as there is so little literature concerning starling welfare. One factor that has been investigated is the effect of lighting quality on both starling welfare and behaviour (Maddocks

et al. 2002a–c, Greenwood *et al.* 2004, Evans *et al.* 2006). Less than 10% of articles mentioned details of lighting environment aside from lighting schedules. This suggests that researchers may not be aware of the particular welfare needs of their study species, and raises concerns about the reporting of husbandry in published papers.

Although the husbandry standards reviewed in this article did fall below new European guidelines, it is not surprising given these guidelines arise from the first document to relate directly to starling welfare published in 2001 (Hawkins *et al.* 2001). Perhaps, a more surprising finding was the amount of methodological detail relating to the birds and their husbandry that could not be obtained from the articles. Basic information that was consistently not reported included: only half of the articles reported the sex of birds, only 41 articles reported temperature and only 33 articles reported the fate of birds. Journals differ in the amount of methodological detail required (Table 5). Many details not included in guidelines for authors are implicitly required for replication, but as this review demonstrates there is a lack of consistency in the reporting of methodological detail (for similar argument see Wurbel 2007).

We found considerable variance in the husbandry of captive starlings between

institutions. Enclosure volumes, for instance, varied between 0.02 and 720 m³, aspect ratios varied from 0.31–10.20 and 12 different lighting schedules were employed from continuous lighting to 8 h of light. Minimum standards must be met to safeguard the welfare of experimental animals, for instance the cage mentioned above with a volume of 0.02 m³ (dimensions: 0.36 × 0.23 × 0.28 m) is clearly unacceptable for housing starlings since it is smaller than a starling’s wingspan. However, optimal conditions will vary depending on the experiment being conducted. For example, in the case of cage size, the largest cages may not be preferable (for either birds or experimenters) if birds need to be caught on a regular basis. Within acceptable ranges, some degree of variance may actually be desirable to reveal interesting and unforeseen effects. For example, Bateson (2002) found small uncontrolled variations in temperature accounted for individual variation in foraging decisions.

It is clear from this review that current practice in laboratory starling husbandry is far from meeting new European guidelines. For example, the majority of articles used cages below the new minimum size recommended. It is vital that more research be conducted into the welfare of starlings, and other less commonly used species, to inform

Table 5 Information required by journal’s guidelines to authors

	Animal behaviour	General and comparative endocrinology	Hormones and behaviour
Sex	✓		
Origin	✓		✓
Time in captivity			✓
Procedures – non-invasive			
Procedures – invasive	?		✓
Fate	?		
Housing type	✓		
Housing dimensions	✓		
Stocking density	✓		
Enrichment	✓		
Food/Water – access	?		
Food/Water – other details	✓		
Lighting – schedule	✓		
Lighting – other details			
Temperature	✓		
Legislation/Guidelines	?	✓	?

✓ = requirement, ? = possible requirement

future guidelines and legislation, and thus maximize the welfare of captive animals.

Acknowledgements This study was funded by a BBSRC Committee Studentship to LA and a Royal Society University Research Fellowship to MB. We thank Stephanie Matheson, Innes Cuthill and an anonymous referee for helpful comments on the manuscript.

References

- Absil P, Pinxten R, Balthazart J, Eens M (2003a) Effect of age and testosterone on autumnal neurogenesis in male European starlings (*Sturnus vulgaris*). *Behavioural Brain Research* **143**, 15–30
- Absil P, Pinxten R, Balthazart J, Eens M (2003b) Effects of testosterone on reelin expression in the brain of male European starlings. *Cell and Tissue Research* **312**, 81–93
- Atasever A, Gumussoy KS (2004) Pathological, clinical and mycological findings in experimental aspergillosis infections of starlings. *Journal of Veterinary Medicine. A, Physiology, Pathology, Clinical Medicine* **51**, 19–22
- Auger CJ, Bentley GE, Auger AP, Ramamurthy M, Ball GF (2002) Expression of camp response element binding protein-binding protein in the song control system and hypothalamus of adult European starlings (*Sturnus vulgaris*). *Journal of Neuroendocrinology* **14**, 805–13
- Bateson M (2002) Context-dependent foraging choices in risk-sensitive starlings. *Animal Behaviour* **64**, 251–60
- Bateson M, Kacelnik A (1995) Preferences for fixed and variable food sources: Variability in amount and delay. *Journal of the Experimental Analysis of Behavior* **63**, 313–29
- Bautista LM, Tinbergen J, Kacelnik A (2001) To walk or to fly? How birds choose among foraging modes. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 1089–94
- Bee MA, Klump GM (2004) Primitive auditory stream segregation: a neurophysiological study in the songbird forebrain. *Journal of Neurophysiology* **92**, 1088–104
- Bentley GE (2003) Melatonin receptor density in area x of European starlings is correlated with reproductive state and is unaffected by plasma melatonin concentration. *General and Comparative Endocrinology* **134**, 187–92
- Bentley GE, Dawson A, Goldsmith AR (2000) Lack of gonadotrophin-releasing hormone (GnRH) neuron response to decreasing photoperiod in thyroidectomized male starlings (*Sturnus vulgaris*). *Journal of Experimental Zoology* **287**, 74–9
- Blackwell BF, Bernhardt GE, Dolbeer RA (2002) Lasers as non-lethal avian repellents. *Journal of Wildlife Management* **66**, 250–8
- Braaten RF (2000) Multiple levels of representation of song by European starlings (*Sturnus vulgaris*): open-ended categorization of starling song types and differential forgetting of song categories and exemplars. *Journal of Comparative Psychology* **114**, 61–72
- Buchanan KL, Spencer KA, Goldsmith AR, Catchpole CK (2003) Song as an honest signal of past developmental stress in the European starling (*Sturnus vulgaris*). *Proceedings of the Royal Society of London Series B—Biological Sciences* **270**, 1149–56
- Butler KL, Fitzgerald SD, Berry DE, *et al.* (2001) Experimental inoculation of European starlings (*Sturnus vulgaris*) and American crows (*Corvus brachyrhynchos*) with *Mycobacterium bovis*. *Avian Diseases* **45**, 709–18
- Clark L, Bryant B, Mezine I (2000) Bird aversive properties of methyl anthranilate, yucca, xanthoxylum, and their mixtures. *Journal of Chemical Ecology* **26**, 1219–34
- Clubb R, Mason G (2003) Captivity effects on wide-ranging carnivores. *Nature* **425**, 473
- Congiu L, Chicca M, Pilaastro A, Turchetto M, Tallandini L (2000) Effects of chronic dietary cadmium on hepatic glutathione levels and glutathione peroxidase activity in starling (*Sturnus vulgaris*). *Archives of Environmental Contamination and Toxicology* **38**, 357–61
- Council of Europe (1986) *European Convention for the Protection of Vertebrate Animals used for Experimental and other Scientific Purposes (ETS 123)*. Strasbourg: Council of Europe
- Cousillas H, Richard J, Mathelier M, *et al.* (2004) Experience-dependent neuronal specialization and functional organization in the central auditory area of a songbird. *European Journal of Neuroscience* **19**, 3343
- Cuthill IC, Hindmarsh A (1985) Increase in starling song activity with removal of mate. *Animal Behaviour* **33**, 326–7
- Cuthill IC, Maddocks SA, Weall CV, Jones EKM (2000) Body mass regulation in response to changes in feeding predictability and overnight energy expenditure. *Behavioral Ecology* **11**, 189–95
- Dawson A (2001) The effects of a single long photoperiod on induction and dissipation of reproductive photorefractoriness in European starlings. *General and Comparative Endocrinology* **121**, 316–24
- Dawson A (2003a) A comparison of the annual cycles in testicular size and moult in captive European starlings, *Sturnus vulgaris* during their first and second years. *Journal of Avian Biology* **34**, 119–23
- Dawson A (2003b) A detailed analysis of primary feather moult in the common starling *Sturnus vulgaris* – new feather mass increases at a constant rate. *Ibis* **145**, E69–76

- Dawson A (2004a) The effects of delaying the start of moult on the duration of moult, primary feather growth rates and feather mass in common starlings *Sturnus vulgaris*. *Ibis* **146**, 493–500
- Dawson A (2004b) Evidence against a period of relative photorefractoriness during the recovery of photosensitivity in common starlings. *General and Comparative Endocrinology* **136**, 117–21
- Dawson A, Van't Hof TJ (2002) Ontogeny of the daily profile of plasma melatonin in European starlings raised under long or short photoperiods. *Journal of Biological Rhythms* **17**, 259–65
- Dawson A, Hinsley SA, Ferns PN, Bonser RH, Eccleston L (2000) Rate of moult affects feather quality: A mechanism linking current reproductive effort to future survival. *Proceedings of the Royal Society of London Series B—Biological Sciences* **267**, 2093–8
- Dawson A, Talbot RT, Dunn IC, Sharp PJ (2002) Changes in basal hypothalamic chicken gonadotropin-releasing hormone and vasoactive intestinal polypeptide associated with a photo-induced cycle in gonadal maturation and prolactin secretion in intact and thyroidectomized starlings (*Sturnus vulgaris*). *Journal of Neuroendocrinology* **14**, 533–9
- De Cock R, Matthysen E (2001) Do glow-worm larvae (Coleoptera: Lampyridae) use warning coloration? *Ethology* **107**, 1019–33
- De Ridder E, Pinxten R, Mees V, Eens M (2002) Short- and long-term effects of male-like concentrations of testosterone on female European starlings (*Sturnus vulgaris*). *Auk* **119**, 487–97
- Devereux CL, McKeever CU, Benton TG, Whittingham MJ (2004) The effect of sward height and drainage on common starlings *Sturnus vulgaris* and northern lapwings *Vanellus vanellus* foraging in grassland habitats. *Ibis* **146**, 115–22
- Duffy DL, Ball GF (2002) Song predicts immunocompetence in male European starlings (*Sturnus vulgaris*). *Proceedings of the Royal Society of London Series B—Biological Sciences* **269**, 847–52
- Duffy DL, Bentley GE, Drazen DL, Ball GF (2000) Effects of testosterone on cell-mediated and humoral immunity in non-breeding adult European starlings. *Behavioral Ecology* **11**, 654–62
- Eens M, Pinxten R, Verheyen R (1991) Male song as a cue for mate choice in the European starling. *Behavioural Brain Research* **116**, 210–38
- European Commission (2006) *Questionnaire for Experts and Organisations on the Preliminary Findings of an Impact Assessment of Options for the Revision of the Directive for the Protection of Animals used in Experiments*. See http://ec.europa.eu/environment/chemicals/lab_animals/pdf/iv_housing_and_care.pdf (last accessed 1 March 2006)
- European Union (1986) *Council Directive 86/609/EEC On the Approximation of Laws, Regulations and Administrative Provisions of the Member States Regarding the Protection of Animals used for Experimental and Other Scientific Purposes*. Strasbourg: Council of Europe
- Evans JE, Cuthill I, Bennett ATD (2006) The effect of flicker from fluorescent lights on mate choice in captive birds. *Animal Behaviour* **72**, 393–400
- Fawcett TW, Skinner AMJ, Goldsmith AR (2002) A test of imitative learning in starlings using a two-action method with an enhanced ghost control. *Animal Behaviour* **64**, 547–56
- Feare C (1984) *The Starling*. Oxford: Oxford University Press
- Fernández-Juricic E, Kacelnik A (2004) Information transfer and gain in flocks: the effects of quality and quantity of social information at different neighbour distances. *Behavioral Ecology and Sociobiology* **55**, 502–11
- Fernández-Juricic E, Siller S, Kacelnik A (2004) Flock density, social foraging, and scanning: an experiment with starlings. *Behavioral Ecology* **15**, 371–9
- Gentner T, Margoliash D (2003) Neuronal populations and single cells representing learned auditory objects. *Nature* **424**, 669–74
- Gentner TQ, Hulse SH, Duffy D, Ball GF (2001) Response biases in auditory forebrain regions of female songbirds following exposure to sexually relevant variation in male song. *Journal of Neurobiology* **46**, 48–58
- Gentner TQ, Hulse SH (2000a) Female European starling preference and choice for variation in conspecific male song. *Animal Behaviour* **59**, 443–58
- Gentner TQ, Hulse SH (2000b) Perceptual classification based on the component structure of song in European starlings. *Journal of the Acoustical Society of America* **107**, 3369–81
- Gentner TQ, Hulse SH, Ball GF (2004) Functional differences in forebrain auditory regions during learned vocal recognition in songbirds. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural and Behavioral Physiology* **190**, 1001–10
- Gentner TQ, Hulse SH, Bentley GE, Ball GF (2000) Individual vocal recognition and the effect of partial lesions to HVC on discrimination, learning, and categorization of conspecific song in adult songbirds. *Journal of Neurobiology* **42**, 117–33
- George I, Vernier B, Richard JP, Hausberger M, Cousillas H (2004) Hemispheric specialization in the primary auditory area of awake and anesthetized starlings (*Sturnus vulgaris*). *Behavioral Neuroscience* **118**, 597–610
- Gill EL (1994) Environmental enrichment for captive starlings. *Animal Technology* **45**, 89–93

- Gill EL, Chivers RE, Ellis SC, *et al.* (1995) Turf as a means of enriching the environment of captive starlings (*Sturnus vulgaris*). *Animal Technology* **46**, 97–102
- Goslow GE, Wilson D, Poore SO (2000) Neuromuscular correlates to the evolution of flapping flight in birds. *Brain Behavior and Evolution* **55**, 85–99
- Greenwood VJ, Smith EL, Church SC, Partridge JC (2003) Behavioural investigation of polarisation sensitivity in the Japanese quail (*Coturnix coturnix japonica*) and the European starling (*Sturnus vulgaris*). *Journal of Experimental Biology* **206**, 3201–10
- Greenwood VJ, Smith EL, Cuthill IC, *et al.* (2002) Do European starlings prefer light environments containing UV? *Animal Behaviour* **64**, 923–8
- Greenwood VJ, Smith EL, Goldsmith AR, *et al.* (2004) Does the flicker frequency of fluorescent lighting affect the welfare of captive European starlings? *Applied Animal Behaviour Science* **86**, 145–59
- Gwinner H, Van't Hof T, Zeman M (2002) Hormonal and behavioral responses of starlings during a confrontation with males or females at nest boxes during the reproductive season. *Hormones and Behavior* **42**, 21–31
- Hambly C, Pinshow B, Wiersma P, *et al.* (2004) Comparison of the cost of short flights in a nectarivorous and a non-nectarivorous bird. *Journal of Experimental Biology* **207**, 3959–68
- Hausberger M, Leppelsack E, Richard JP, Leppelsack HJ (2000) Neuronal bases of categorization in starling song. *Behavioural Brain Research* **114**, 89–95
- Hawkins P, Bairlein F, Duncan I, *et al.* (2003) *Future Principles for Housing and Care of Laboratory Birds: Report for the Revision of the Council of Europe Convention ETS123 Appendix A for Birds*. Strasbourg: Council of Europe
- Hawkins P, Morton DB, Cameron D, *et al.* (2001) Laboratory birds: refinements in husbandry and procedures. *Laboratory Animals* **35**, S1–163
- Henry L, Hausberger M (2001) Differences in the social context of song production in captive male and female European starlings. *Comptes Rendus de l'Academie des Sciences. Serie III, Sciences de la Vie* **324**, 1167–74
- Hile AG (2004) Avoidance of plant secondary compounds by European starlings: Citronellyls. *Crop Protection* **23**, 973–8
- Hile AG, Shan ZX, Zhang SZ, Block E (2004) Aversion of European starlings (*Sturnus vulgaris*) to garlic oil treated granules: garlic oil as an avian repellent. Garlic oil analysis by nuclear magnetic resonance spectroscopy. *Journal of Agricultural and Food Chemistry* **52**, 2192–6
- Hodgson JC, Spielman A, Komar N, *et al.* (2001) Interrupted blood-feeding by *Culiseta melanura* (Diptera: Culicidae) on European starlings. *Journal of Medical Entomology* **38**, 59–66
- Hofer SB, Klump GM (2003) Within- and across-channel processing in auditory masking: a physiological study in the songbird forebrain. *Journal of Neuroscience* **23**, 5732–9
- Kacelnik A (1984) Central place foraging in starlings (*Sturnus-vulgaris*). 1. Patch residence time. *Journal of Animal Ecology* **53**, 283–99
- Kacelnik A, Marsh B (2002) Cost can increase preference in starlings. *Animal Behaviour* **63**, 245–50
- Kumar V, Gwinner E, Van't Hof TJ (2000) Circadian rhythms of melatonin in European starlings exposed to different lighting conditions: relationship with locomotor and feeding rhythms. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural and Behavioral Physiology* **186**, 205–15
- Langemann U, Klump GM (2001) Signal detection in amplitude-modulated maskers. I. Behavioural auditory thresholds in a songbird. *European Journal of Neuroscience* **13**, 1025–32
- Lima S (1983) Patch sampling behavior of starlings foraging in simple patchy environments. *American Zoologist* **23**, 897
- Lucas JR, Freeberg TM, Krishnan A, Long GR (2002) A comparative study of avian auditory brainstem responses: correlations with phylogeny and vocal complexity, and seasonal effects. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural and Behavioral Physiology* **188**, 981–92
- Maddocks SA, Goldsmith AR, Cuthill IC (2001) The influence of flicker rate on plasma corticosterone levels of European starlings, *Sturnus vulgaris*. *General and Comparative Endocrinology* **124**, 315–20
- Maddocks SA, Bennett ATD, Cuthill IC (2002a) Rapid behavioural adjustments to unfavourable light conditions in European starlings (*Sturnus vulgaris*). *Animal Welfare* **11**, 95–101
- Maddocks SA, Goldsmith AR, Cuthill IC (2002b) Behavioural and physiological effects of absence of ultraviolet wavelengths on European starlings *Sturnus vulgaris*. *Journal of Avian Biology* **33**, 103–6
- Maddocks SA, Bennett ATD, Hunt S, Cuthill IC (2002c) Context-dependent visual preferences in starlings and blue tits: mate choice and light environment. *Animal Behaviour* **63**, 69–75
- Marsh B, Kacelnik A (2002) Framing effects and risky decisions in starlings. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 3352–5
- Marsh B, Schuck-Paim C, Kacelnik A (2004) Energetic state during learning affects foraging choices in starlings. *Behavioral Ecology* **15**, 369–99

- Mountjoy D, Lemon R (1991) Song as an attractant for male and female European starlings, and the influence of song complexity on their response. *Behavioral Ecology and Sociobiology* **28**, 97–100
- National Research Council (1996) *Guide for the Care and Use of Laboratory Animals*. Bethesda: Office of Science and Health Reports
- Nephew BC, Romero LM (2003) Behavioral, physiological, and endocrine responses of starlings to acute increases in density. *Hormones and Behavior* **44**, 222–32
- Nephew BC, Kahn SA, Romero LM (2003) Heart rate and behavior are regulated independently of corticosterone following diverse acute stressors. *General and Comparative Endocrinology* **133**, 173–80
- Nieder A, Klump GM (2001) Signal detection in amplitude-modulated maskers. II. Processing in the songbird's auditory forebrain. *European Journal of Neuroscience* **13**, 1033–44
- Oberweger K, Goller F (2001) The metabolic cost of birdsong production. *Journal of Experimental Biology* **204**, 3379–88
- Olsson O, Brown JS, Smith HG (2001) Gain curves in depletable food patches: a test of five models with European starlings. *Evolutionary Ecology Research* **3**, 285–310
- Olsson O, Bruun M, Smith HG (2002) Starling foraging success in relation to agricultural land-use. *Ecography* **25**, 363–71
- Pilz KM, Quiroga M, Schwabl H, Adkins-Regan E (2004) European starling chicks benefit from high yolk testosterone levels during a drought year. *Hormones and Behavior* **46**, 179–92
- Pinxten R, De Ridder E, Balthazart J, Berghman L, Eens M (2000) The effect of castration on aggression in the nonbreeding season is age-dependent in male European starlings. *Behaviour* **137**, 647–61
- Pinxten R, De Ridder E, Balthazart J, Eens M (2002) Context-dependent effects of castration and testosterone treatment on song in male European starlings. *Hormones and Behavior* **42**, 307–18
- Pinxten R, De Ridder E, De Cock M, Eens M (2003a) Castration does not decrease nonreproductive aggression in yearling male European starlings (*Sturnus vulgaris*). *Hormones and Behavior* **43**, 394–401
- Pinxten R, de Ridder E, Eens M (2003b) Female presence affects male behavior and testosterone levels in the European starling (*Sturnus vulgaris*). *Hormones and Behavior* **44**, 103–9
- Poirier C, Henry L, Mathelier M, et al. (2004) Direct social contacts override auditory information in the song-learning process in starlings (*Sturnus vulgaris*). *Journal of Comparative Psychology* **118**, 179–93
- Poole T, Dawkins M (1999) Environmental enrichment for vertebrates. In: *The UFAW Handbook on the Care and Management of Laboratory Animals* (Poole T, ed). London: Blackwell Science, 13–20
- Powell G (1974) Experimental analysis of social value of flocking by starlings (*Sturnus-vulgaris*) in relation to predation and foraging. *Animal Behaviour* **22**, 501–5
- Rafacz M, Templeton JJ (2003) Environmental unpredictability and the value of social information for foraging starlings. *Ethology* **109**, 951–60
- Rayner JMV, Viscardi PW, Ward S, Speakman JR (2001) Aerodynamics and energetics of intermittent flight in birds. *American Zoologist* **41**, 188–204
- Remage-Healey L, Romero LM (2000) Daily and seasonal variation in response to stress in captive starlings (*Sturnus vulgaris*): glucose. *General and Comparative Endocrinology* **119**, 60–8
- Remage-Healey L, Romero LM (2001) Corticosterone and insulin interact to regulate glucose and triglyceride levels during stress in a bird. *American Journal of Physiology. Regulatory Integrative and Comparative Physiology* **281**, R994–1003
- Remage-Healey L, Romero LM (2002) Corticosterone and insulin interact to regulate plasma glucose but not lipid concentrations in moulting starlings. *General and Comparative Endocrinology* **129**, 88–94
- Riters LV, Alger SJ (2004) Neuroanatomical evidence for indirect connections between the medial preoptic nucleus and the song control system: possible neural substrates for sexually motivated song. *Cell and Tissue Research* **316**, 35–44
- Riters LV, Baillien M, Eens M, et al. (2001) Seasonal variation in androgen-metabolizing enzymes in the diencephalon and telencephalon of the male European starling (*Sturnus vulgaris*). *Journal of Neuroendocrinology* **13**, 985–97
- Riters LV, Eens M, Pinxten R, Ball GF (2002) Seasonal changes in the densities of alpha(2)-noradrenergic receptors are inversely related to changes in testosterone and the volumes of song control nuclei in male European starlings. *Journal of Comparative Neurology* **444**, 63–74
- Riters LV, Eens M, Pinxten R, et al. (2000) Seasonal changes in courtship song and the medial preoptic area in male European starlings (*Sturnus vulgaris*). *Hormones and Behavior* **38**, 250–61
- Riters LV, Teague DP (2003) The volumes of song control nuclei, HVC and IMAN, relate to differential behavioral responses of female European starlings to male songs produced within and outside of the breeding season. *Brain Research* **978**, 91–8
- Romero LM, Remage-Healey L (2000) Daily and seasonal variation in response to stress in captive starlings (*Sturnus vulgaris*): corticosterone. *General and Comparative Endocrinology* **119**, 52–9
- Rothery P, Wyllie I, Newton I, Dawson A, Osborn D (2001) The timing and duration of moult in adult

- starlings *Sturnus vulgaris* in East-central England. *Ibis* **143**, 435–41
- Sayre RW, Clark L (2001) Effect of primary and secondary repellents on European starlings: an initial assessment. *Journal of Wildlife Management* **65**, 461–9
- Schuck-Paim C, Kacelnik A (2002) Rationality in risk-sensitive foraging choices by starlings. *Animal Behaviour* **64**, 869–79
- Schuck-Paim C, Pompilio L, Kacelnik A (2004) State-dependent decisions cause apparent violations of rationality in animal choice. *PLoS Biology* **2**, e402
- Seibels B, Lamberski N, Gregory CR, Slifka K, Hagerman AE (2003) Effective use of tea to limit dietary iron available to starlings (*Sturnus vulgaris*). *Journal of Zoo and Wildlife Medicine* **34**, 314–16
- Sockman KW, Gentner TQ, Ball GF (2002) Recent experience modulates forebrain gene-expression in response to mate-choice cues in European starlings. *Proceedings of the Royal Society of London Series B—Biological Sciences* **269**, 2479–85
- Sockman KW, Williams TD, Dawson A, Ball GF (2004) Prior experience with photostimulation enhances photo-induced reproductive development in female European starlings: a possible basis for the age-related increase in avian reproductive performance. *Biology of Reproduction* **71**, 979–86
- Staples J, Krall B, Bartelt R, Whitman D (2002) Chemical defense in the plant bug *Lopidea robiniae* (Uhler). *Journal of Chemical Ecology* **28**, 601–15
- Swaddle JP, Lockwood R (2003) Wingtip shape and flight performance in the European starling *Sturnus vulgaris*. *Ibis* **145**, 457–64
- Swaddle JP, Pruett-Jones S (2001) Starlings can categorize symmetry differences in dot displays. *American Naturalist* **158**, 300–7
- Swaddle JP, Ruff DA (2004) Starlings have difficulty in detecting dot symmetry: implications for studying fluctuating asymmetry. *Behaviour* **141**, 29–40
- Swaddle JP, Che JPK, Clelland RE (2004) Symmetry preference as a cognitive by-product in starlings. *Behaviour* **141**, 469–78
- Talling JC, Inglis IR, Van Driel KS, Young J, Giles S (2002) Effect of hunger on starlings' preferences for food sources associated with variability or uncertainty. *Behaviour* **139**, 1223–35
- Templeton JJ, Gonzalez DP (2004) Reverse lateralization of visual discriminative abilities in the European starling. *Animal Behaviour* **67**, 783–8
- Tinbergen J (1981) Foraging decisions in starlings (*Sturnus-vulgaris* L). *Ardea* **69**, 1–67
- Van Meir V, Verhoye M, Absil P, et al. (2004) Differential effects of testosterone on neuronal populations and their connections in a sensorimotor brain nucleus controlling song production in songbirds: a manganese enhanced-magnetic resonance imaging study. *Neuroimage* **21**, 914–23
- Vasquez RA, Kacelnik A (2000) Foraging rate versus sociality in the starling *Sturnus vulgaris*. *Proceedings of the Royal Society of London Series B—Biological Sciences* **267**, 157–64
- Výoh P, Zeman M, Buyse J, Jurani M, Decuypere E (2001) Developmental changes in thyrotropic and somatotropic effect of TRR in precocial Japanese quail and altricial European starlings. *General and Comparative Endocrinology* **124**, 293–9
- Ward S, Moller U, Rayner JMV, et al. (2001) Metabolic power, mechanical power and efficiency during wind tunnel flight by the European starling *Sturnus vulgaris*. *Journal of Experimental Biology* **204**, 3311–22
- Ward S, Möller U, Rayner JMV, Jackson DM, Nachtigall W, Speakman JR (2004) Metabolic power of European starlings *Sturnus vulgaris* during flight in a wind tunnel, estimated from heat transfer modelling, doubly labelled water and mask respirometry. *Journal of Experimental Biology* **207**, 4291–8
- West M, King A (1990) Mozart starling. *American Scientist* **78**, 106–14
- Williams EV, Swaddle JP (2003) Moulting, flight performance and wingbeat kinematics during take-off in European starlings *Sturnus vulgaris*. *Journal of Avian Biology* **34**, 371–8
- Wurbel H (2007) Publications should include an animal-welfare section. *Nature* **446**, 257
- Young KA, Ball GF, Nelson RJ (2001) Photoperiod-induced testicular apoptosis in European starlings (*Sturnus vulgaris*). *Biology of Reproduction* **64**, 706–13